

Pediatric and Adolescent Immunization Schedule

Vaccines are listed under the recommended administration in the dark **PINK** shaded boxes.

The **GREY** shaded boxes indicate the administration ages if previous doses were missed.

	BIRTH	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS	2-3 YEARS	4-6 YEARS	7-10 YEARS	11-12 YEARS	13- 18 YEARS
Hepatitis B (HepB)	HepB	HepB			HepB									
Rotavirus (RV)			RV	RV	RV ¹									
Diphtheria, Tetanus, Pertussis (DTaP)			DTaP	DTaP	DTaP	Note ²	DTaP				DTaP		Tdap	
H. Influenzae type b (Hib)			Hib	Hib	Hib ³	Hib								
Pneumococcal (PCV)			PCV	PCV	PCV	PCV								
Inactivated Polio Virus (IPV)			IPV	IPV	IPV						IPV			
Influenza					Influenza (Yearly)									
Measles, Mumps, Rubella (MMR)						MMR			Note ⁴		MMR			
Varicella (Chicken Pox)						Varicella			Note ⁵		Varicella			
Hepatitis A (HepA)						HepA (two doses) ⁶								
Meningococcal (MCV)													MCV	
Human Papillomavirus (HPV)													HPV (three doses) ⁷	

1. If Rotarix® is administered at 2 and 4 months, 6 month dose is not indicated.

2. The fourth dose of DTaP may be administered as early as age 12 months, provided 6 months have elapsed since the third dose.

3. Six month dose not required if PRP-OMP is used at ages 2 and 4 months.

4. The second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.

5. The second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.

6. Administer 2 doses at least 6 months apart.

7. Administer the second dose 1-2 months after the first dose and the third dose 6 months after the first dose.